arXiv:1702.06926v1 [math.GR] 22 Feb 2017

*International Journal of Group Theory Vol.* XX *No. X (201X), pp XX-XX.*

# ON VARIETAL CAPABILITY OF INFINITE DIRECT PRODUCTS OF GROUPS

HANIEH MIREBRAHIMI* AND BEHROOZ MASHAYEKHY

Communicated by

Abstract. Recently, the authors gave some conditions under which a direct product of finitely many groups is $\mathcal{V}-$capable if and only if each of its factors is $\mathcal{V}-$capable for some varieties $\mathcal{V}$. In this paper, we extend this fact to any infinite direct product of groups. Moreover, we conclude some results for $\mathcal{V}-$capability of direct products of infinitely many groups in varieties of abelian, nilpotent and polynilpotent groups.

## 1. Introduction

R. Baer [1] initiated an investigation of the question "which conditions a group $G$ must fulfill in order to be the group of inner automorphisms of a group $E$?", that is $G \cong E/Z(E)$. Following M. Hall and J. K. Senior [5], such a group $G$ is called *capable*. Baer [1] determined all capable groups which are direct sums of cyclic groups. As P. Hall [4] mentioned, characterizations of capable groups are important in classifying groups of prime-power order.

F. R. Beyl, U. Felgner and P. Schmid [2] proved that every group $G$ possesses a uniquely determined central subgroup $Z^*(G)$ which is minimal subject to being the image in $G$ of the center of some central

MSC(2010): Primary: 20E10; Secondary: , 20K25, 20E34, 20D15, 20F18.
Keywords: Capable group, Direct product, Variety of groups, $\mathcal{V}-$capable group, direct limit.
Received: 30 April 2009, Accepted: 21 June 2010.
*Corresponding author
© 2011 University of Isfahan.

extension of $G$. This $Z^*(G)$ is characteristic in $G$ and is the image of the center of every stem cover of $G$. Moreover, $Z^*(G)$ is the smallest central subgroup of $G$ whose factor group is capable [2]. Hence $G$ is capable if and only if $Z^*(G) = 1$. They showed that the class of all capable groups is closed under the direct products. Also, they presented a condition in which the capability of a direct product of finitely many of groups implies the capability of each of the factors. Moreover, they proved that if $N$ is a central subgroup of $G$, then $N \subseteq Z^*(G)$ if and only if the mapping $M(G) \to M(G/N)$ induced by the natural epimorphism, is monomorphism.

Then M. R. R. Moghadam and S. Kayvanfar [10] generalized the concept of capability to $\mathcal{V}-$capability for a group $G$. They introduced the subgroup $(V^*)^*(G)$ which is associated with the variety $\mathcal{V}$ defined by a set of laws $V$ and a group $G$ in order to establish a necessary and sufficient condition under which $G$ can be $\mathcal{V}-$capable. They also showed that the class of all $\mathcal{V}-$capable groups is closed under the direct products. Moreover, they exhibited a close relationship between the groups $\mathcal{V}M(G)$ and $\mathcal{V}M(G/N)$, where $N$ is a normal subgroup contained in the marginal subgroup of $G$ with respect to the variety $\mathcal{V}$. Using this relationship, they gave a necessary and sufficient condition for a group $G$ to be $\mathcal{V}-$capable.

The authors [7] presented some conditions in which the $\mathcal{V}-$capablity of a direct product of finitely many groups implies the $\mathcal{V}-$capablity of each of its factors. In this paper, we extend this fact to direct product of an infinite family of groups. Also, we deduce some new results about the $\mathcal{V}-$capability of direct product of infinitely many groups, where $\mathcal{V}$ is the variety of abelian, nilpotent, or polynilpotent groups.

## 2. **Main Results**

Suppose that $\mathcal{V}$ is a variety of groups defined by the set of laws $V$. A group $G$ is said to be $\mathcal{V}-$capable if there exists a group $E$ such that $G \cong E/V^*(E)$, where $V^*(E)$ is the marginal subgroup of $E$, which is defined as follows [6]:

$$\{g \in E \mid v(x_1, x_2, ..., x_n) = v(x_1, x_2, ..., gx_i, x_{i+1}, ..., x_n)$$
$$\forall x_1, x_2, ..., x_n \in E,\ \forall i \in \{1, 2, ..., n\}\}.$$

If $\psi : E \to G$ is a surjective homomorphism with $ker\psi \subseteq V^*(E)$, then the intersection of all subgroups of the form $\psi(V^*(E))$ is denoted by

$(V^*)^*(G)$. It is obvious that $(V^*)^*(G)$ is a characteristic subgroup of $G$ contained in $V^*(G)$. If $\mathcal{V}$ is the variety of abelian groups, then the subgroup $(V^*)^*(G)$ is the same as $Z^*(G)$ and in this case $\mathcal{V}-$capability is equal to capability [10]. In the following, there are some results which we need them in sequel.

**Theorem 2.1.** [10] *(i) A group $G$ is $\mathcal{V}-$capable if and only if $(V^*)^*(G) = 1$.*
*(ii) If $\{G_i\}_{i\in I}$ is a family of groups, then $(V^*)^*(\prod_{i\in I} G_i) \subseteq \prod_{i\in I}(V^*)^*(G_i)$.*

As a consequence, if the $G_i$'s are $\mathcal{V}-$capable groups, then $G = \prod_{i\in I} G_i$ is also $\mathcal{V}-$capable. In the above theorem, the equality does not hold in general (see Example 2.3 of [7]).

**Theorem 2.2.** [10] *Let $\mathcal{V}$ be a variety of groups with a set of laws $V$. Let $G$ be a group and $N$ be a normal subgroup with the property $N \subseteq V^*(G)$. Then $N \subseteq (V^*)^*(G)$ if and only if the homomorphism induced by the natural map $\mathcal{V}M(G) \to \mathcal{V}M(G/N)$ is a monomorphism.*

We recall that the Baer-invariant of a group $G$, with the free presentation $F/R$, with respect to the variety $\mathcal{V}$, denoted by $\mathcal{V}M(G)$, is

$$\mathcal{V}M(G) = \frac{R \cap V(F)}{[RV^*F]} \ ,$$

where $V(F)$ is the verbal subgroup of $F$ with respect to $\mathcal{V}$ and

$$[RV^*F] =< v(f_1, \ldots, f_{i-1}, f_i r, f_{i+1}, \ldots, f_n)v(f_1, \ldots, f_i, \ldots, f_n)^{-1} | r \in R,$$

$$f_i \in F, v \in V, 1 \leq i \leq n, n \in \mathbf{N} > \ .$$

It is known that the Baer-invariant of a group $G$ is always abelian and independent of the choice of the presentation of $G$. Also if $\mathcal{V}$ is the variety of abelian groups, then the Baer-invariant of $G$ will be $R \cap F'/[R, F] \cong M(G)$, where $M(G)$ is the Schur multiplier of $G$ (see [6]).

**Theorem 2.3.** [7] *Let $\mathcal{V}$ be a variety, $A$ and $B$ be two groups with $\mathcal{V}M(A \times B) \cong \mathcal{V}M(A) \times \mathcal{V}M(B)$, then $(V^*)^*(A \times B) = (V^*)^*(A) \times (V^*)^*(B)$. Consequently, $A \times B$ is $\mathcal{V}$-capable if and only if $A$ and $B$ are both $\mathcal{V}$-capable.*

**Theorem 2.4.** [8]*Let $\{G_i; \phi_i^j, I\}$ be a directed system of groups. Then, for a given variety $\mathcal{V}$, the Baer-invariant preserves direct limit, that is $\mathcal{V}M(\varinjlim G_i) = \varinjlim \mathcal{V}M(G_i)$.*

**Lemma 2.5.** *For any family of groups $\{G_i\}_{i\in I}$, consider the directed system $\{\mathcal{G}_{I_\lambda}, \phi_\lambda^{\lambda'}, \Lambda\}$ consisting of all finite direct products $\mathcal{G}_{I_\lambda} = \prod_{i\in I_\lambda} G_i$ ($I_\lambda$ is a finite subset of $I$), with the natural embedding homomorphisms $\phi_\lambda^{\lambda'} : \mathcal{G}_{I_\lambda} \to \mathcal{G}_{I_{\lambda'}}$ ($I_\lambda \subseteq I_{\lambda'}$). Also, the index set $\Lambda$ is ordered in a directed way so that for any $\lambda, \lambda' \in \Lambda$, $\lambda \leq \lambda'$ if and only if $I_\lambda \subseteq I_{\lambda'}$. Then the direct product $\mathcal{G}_I = \prod_{i\in I} G_i$ is a direct limit of this directed system.*

*Proof.* Let $\mathcal{G} = \varinjlim \mathcal{G}_{I_\lambda}$ be a direct limit of this directed system, with homomorphisms $\phi_\lambda : \mathcal{G}_{I_\lambda} \to \mathcal{G}$. Also, for any $\lambda \in \Lambda$, consider the embedding homomorphism $\tau_\lambda : \mathcal{G}_{I_\lambda} \to \mathcal{G}_I$. Clearly, for any $\lambda, \lambda' \in \Lambda$ with $\lambda \leq \lambda'$, $\tau_{\lambda'}\tau_\lambda^{\lambda'} = \tau_\lambda$. Now, by universal property of $\mathcal{G}$, there exists a unique homomorphism $\phi : \mathcal{G} \to \mathcal{G}_I$ such that for any $\lambda \in \Lambda$, $\phi\phi_\lambda = \tau_\lambda$. To define the inverse homomorphism $\tau : \mathcal{G}_I \to \mathcal{G}$, recall that for any $x = \{x_i\}_{i\in I} \in \mathcal{G}_I$, there exists a finite subset $I_\lambda$ of $I$ that for any $i \in I\backslash I_\lambda$, $x_i$ is trivial in $G_i$. Hence we can consider $x$ as an element of $\mathcal{G}_{I_\lambda}$ and define $\tau(x) = \phi_\lambda(x)$. It is easy to see that for any $\lambda \in \Lambda$, $\tau\tau_\lambda = \phi_\lambda$. Finally, we see that for any $x \in \mathcal{G}_I$, $\phi\tau(x) = \phi\phi_\lambda(x)$, for some $\lambda \in \Lambda$; and so $\phi\tau(x) = \tau_\lambda(x) = x$. Conversely, the equation $\tau\phi = id_{\mathcal{G}}$ holds because of the universal property of the direct limit $\mathcal{G}$. □

By the above notations, we conclude that $\prod_{i\in I} G_i$, $\prod_{i\in I} V^{**}(G_i)$, and $\prod_{i\in I} G_i/V^{**}(G_i)$ are direct limits of directed systems $\{\prod_{i\in I_\lambda} G_i, \phi_\lambda^{\lambda'}, \Lambda\}$, $\{\prod_{i\in I_\lambda} V^{**}(G_i), \bar{\phi}_\lambda^{\lambda'}, \Lambda\}$, and $\{\prod_{i\in I_\lambda} G_i/V^{**}(G_i), \psi_\lambda^{\lambda'}, \Lambda\}$ respectively, where $\bar{\phi}_\lambda^{\lambda'}$'s are restrictions of $\phi_\lambda^{\lambda'}$'s and $\psi_\lambda^{\lambda'}$'s are quotient homomorphisms induced by $\phi_\lambda^{\lambda'}$'s.

Now, suppose that $\{G_i\}_{i\in I}$ is a family of groups in which for any $G_i$ and $G_j$ $(i, j \in I)$, $\mathcal{V}M(G_i \times G_j) \cong \mathcal{V}M(G_i) \times \mathcal{V}M(G_j)$. By Theorem 2.3, $\prod_{i\in I_\lambda}(V^*)^*(G_i) \subseteq (V^*)^*(\prod_{i\in I_\lambda} G_i)$, for any finite subset $I_\lambda$ of $I$. Thus, using Theorem 2.2, we have the following monomorphism

$$\mathcal{V}M(\prod_{i\in I_\lambda} G_i) \hookrightarrow \mathcal{V}M(\frac{\prod_{i\in I_\lambda} G_i}{\prod_{i\in I_\lambda}(V^*)^*(G_i)}).$$

By the fact that direct limit of a directed system preserves exactness of a sequence [8], we obtain the following monomorphism

$$\varinjlim \mathcal{V}M(\prod_{i\in I_\lambda} G_i) \hookrightarrow \varinjlim \mathcal{V}M(\frac{\prod_{i\in I_\lambda} G_i}{\prod_{i\in I_\lambda}(V^*)^*(G_i)}).$$

Using Theorem 2.4, we conclude the monomorphism

$$\mathcal{V}M(\varinjlim \prod_{i\in I_\lambda} G_i) \hookrightarrow \mathcal{V}M(\varinjlim \frac{\prod_{i\in I_\lambda} G_i}{\prod_{i\in I_\lambda}(V^*)^*(G_i)}),$$

and so we have the monomorphism

$$\mathcal{V}M(\prod_{i\in I} G_i) \hookrightarrow \mathcal{V}M(\frac{\prod_{i\in I} G_i}{\prod_{i\in I}(V^*)^*(G_i)}).$$

Finally, by Theorem 2.2, we conclude that

$$\prod_{i\in I}(V^*)^*(G_i) \subseteq (V^*)^*(\prod_{i\in I} G_i).$$

Using these notes, we deduce the following theorem.

**Theorem 2.6.** *Let $\mathcal{V}$ be a variety, $\{G_i\}_{i\in I}$ be a family of groups such that for any $i, j \in I$, $\mathcal{V}M(G_i \times G_j) \cong \mathcal{V}M(G_i) \times \mathcal{V}M(G_j)$. Then $(V^*)^*(\prod_{i\in I} G_i) = \prod_{i\in I}(V^*)^*(G_i)$. Consequently, $\prod_{i\in I} G_i$ is $\mathcal{V}$-capable if and only if each $G_i$ is $\mathcal{V}$-capable.*

**Remark 2.7.** *(i) In the above theorem, the sufficient condition*

$$\mathcal{V}M(A \times B) \cong \mathcal{V}M(A) \times \mathcal{V}M(B)$$

*is not necessary (see Example 2.3$(iii)$ of [7]). Also, this condition is essential and can not be omitted (see Example 2.3$(i), (ii)$ of [7]).*
*(ii) It is known that for varieties of abelian and nilpotent groups, and for any groups $A$ and $B$, $\mathcal{V}M(A \times B) \cong \mathcal{V}M(A) \times \mathcal{V}M(B) \times T$, where $T$ is an abelian group whose elements are tensor products of the elements of $A^{ab}$ and $B^{ab}$ [3], [9]. Hence in these known varieties, the isomorphism $\mathcal{V}M(A \times B) \cong \mathcal{V}M(A) \times \mathcal{V}M(B)$ holds, where both $A^{ab}$ and $B^{ab}$ have finite exponent with $(exp(A^{ab}), exp(B^{ab})) = 1$.*

In the following, using the main theorem and the above remark, we deduce some corollaries which are generalizations of some results of [7] (Remark 2.4$(ii)$, Corollary 2.5 and Example 2.2).

**Corollary 2.8.** *Let $\{G_i\}_{i\in I}$ be a family of groups whose abelianizations have mutually coprime exponents. Then $\prod_{i\in I} G_i$ is capable ($\mathcal{N}_c$-capable) if and only if each $G_i$ is capable ($\mathcal{N}_c$-capable).*

**Corollary 2.9.** *Suppose that $\{G_i\}_{i\in I}$ is a family of groups whose abelianizations have mutually coprime exponents. If $\prod_{i\in I} G_i$ is nilpotent of class at most $c_1$, then it is $\mathcal{N}_{c_1,\cdots,c_s}$-capable if and only if every $G_i$ is $\mathcal{N}_{c_1,\cdots,c_s}$-capable.*

**Corollary 2.10.** *If $\{G_i\}_{i\in I}$ is a family of perfect groups, then $\prod_{i\in I} G_i$ is $\mathcal{V}$-capable if and only if each $G_i$ is $\mathcal{V}$-capable, where $\mathcal{V}$ may be each of these three varieties, 1.variety of abelian groups, 2.variety of nilpotent groups, 3.variety of polynilpotent groups.*

## References


[1] R. Baer, Groups with preassigned central and central quotient groups, *Trans. Amer. Math. Soc.*, (1938), no. 44, 378-412.

[2] F. R. Beyl, U. Felgner, P. Schmid, On groups occurring as center factor groups, *J. Algebra*, (1979), no. 61, 161-177.

[3] G. Ellis, On groups with a finite nilpotent upper central quotient, *Arch. Math.*, (1998), no. 70, 89-96.

[4] P. Hall, The classification of prime-power groups, *J. reine angew. Math.*, (1940), no. 182, 130-141.

[5] M. Hall, Jr., J. K. Senior, *The Groups of Order* $2^n (n \leq 6)$, Macmillan, New York, 1964.

[6] G. Karpilovsky, *The Schur Multiplier*, London Math. Soc. Monographs, New Series 2, Clarendon Press, Oxford University Press, Oxford, 1987.

[7] H. Mirebrahimi, B. Mashayekhy, On varietal capability of direct product of groups, *J. Adv. Res. Pure Math.*, to appear.

[8] M. R. R. Moghaddam, The Baer-invarient and the direct limit, *Mh. Math.*, (1980), no. 90, 37-43.

[9] M. R. R. Moghaddam, The Baer-invarient of a direct product, *Arch. Math.*, (1980), no. 33, 504-511.

[10] M. R. R. Moghaddam, S. Keyvanfar, A new notion derived from varieties of groups. *Algebra Colloq.*, (1997), no. 4:1, 1-11.

[11] D. J. S. Robinson, *A Course in the Theory of Groups*, Second Edition, Springer-Verlag, 1996. mer. Math. Soc., Providence, RI, 2010.



**Hanieh Mirebrahimi**
Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures, Ferdowsi University of Mashhad, P. O. Box 1159-91775, Mashhad, Iran.
Email: h_mirebrahimi@um.ac.ir

**Behrooz Mashayekhy**
Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures, Ferdowsi University of Mashhad, P. O. Box 1159-91775, Mashhad, Iran.
Email: bmashf@um.ac.ir